

NBI-HE-92-12
February, 1992

Hamiltonian Reduction and Classical Extended Superconformal Algebras

Katsushi Ito and Jens Ole Madsen

*The Niels Bohr Institute
University of Copenhagen
Blegdamsvej 17, DK-2100
Copenhagen Ø, Denmark*

Abstract

We present a systematic construction of classical extended superconformal algebras from the hamiltonian reduction of a class of affine Lie superalgebras, which include an even subalgebra $sl(2)$. In particular, we obtain the doubly extended $N = 4$ superconformal algebra $\tilde{A}_\gamma$ from the hamiltonian reduction of the exceptional Lie superalgebra $D(2|1; \gamma/(1 - \gamma))$. We also find the Miura transformation for these algebras and give the free field representation. A W -algebraic generalization is discussed.

In the last few years it has been known that various higher-spin extension of the Virasoro algebra including Zamolodchikov's W algebra[1] can be constructed using the Drinfeld-Sokolov type hamiltonian reduction of affine Lie algebras (or the associated integrable systems) [2],[3] and its generalization [4]-[7]. Supersymmetric extension of the Virasoro algebra is important from the viewpoint of application to superstrings and topological field theory. It has been shown that $N = 1$ and $N = 2$ super W algebras can be obtained from the hamiltonian reduction (or related super Toda field theories) of a special class of affine Lie superalgebras which have the purely odd simple root system [8].

The purpose of this paper is to present a systematic construction of the classical extended superconformal algebra (ESA) from the hamiltonian reduction viewpoint. ESAs with $u(N)$ and $so(N)$ Kac-Moody symmetries, which have been constructed by Knizhnik and Bershadsky [9], have in general non-linear properties similar to those of the W -algebra. Some previous works describe the relation between $N = 1$ and 2 superconformal algebras and Lie superalgebras $osp(N|2)$ [10]. In particular Mathieu has shown that the hamiltonian structure of the $osp(N|2)$ KdV equations leads to the classical $so(N)$ ESA [11], although in his formulation the Lie superalgebraic structure is not obvious.

In the present paper we shall discuss a class of Lie superalgebras which include even subalgebras $\mathfrak{g} \oplus sl(2)$, where $\mathfrak{g}$ is a semi-simple Lie algebra. Using Kac's notation [12], such basic classical Lie superalgebras are classified as follows (see Table 1): $A(n|1)$ ($n \geq 1$), $A(1|0)$ $B(n|1)$, $D(n|1)$, $B(1|n)$, $D(2|1; \alpha)$, $F(4)$ and $G(3)$. By considering the hamiltonian reduction of an even subalgebra $sl(2)$, we will find ESAs with $A_n \oplus u(1)$ (A_1 for $n = 1$), $u(1)$, B_n , D_n , C_n , $A_1 \oplus A_1$, $spin(7)$, G_2 Kac-Moody symmetry¹. In particular for $D(2|1; \alpha)$ the corresponding ESA is shown to be the $N = 4$ doubly extended superconformal algebra [14]. We will also show that the Miura transformation can be naturally obtained by connecting the Drinfeld-Sokolov gauge and the "diagonal" gauge. The present classical

¹For $B(1|n)$ the corresponding algebra has spin 0, 1 and 2 fermionic currents with C_n Kac-Moody symmetry.

formulation is a crucial step for the study of the quantum ESA using the Coulomb gas representation. We note that a bosonic analogue of $u(N)$ ESA has been discussed in refs. [7] by considering the hamiltonian reduction of $sl(N+2)$. However, for other Lie superalgebras there is no bosonic counterpart, especially for $D(2|1; \alpha)$.

We begin by explaining the hamiltonian reduction by taking an affine Lie superalgebra $\hat{\mathbf{g}} = A(n|1)^{(1)}$ at level k associated with a basic classical Lie superalgebra $\mathbf{g} = A(n|1) = sl(N|2)$ ($N = n+1$) as an example. The algebra $\hat{\mathbf{g}}$ consists of an element $(X(z), x_0)$ of the form

$$X(z) = \left(\begin{array}{c|c} X_1(z) & \xi(z) \\ \hline \eta(z) & X_2(z) \end{array} \right), \quad \text{str} X = \text{tr} X_1 - \text{tr} X_2 = 0, \quad (1)$$

where X_1 is a $N \times N$ matrix and X_2 is a 2×2 matrix, whose components are commuting numbers. $\xi = (\xi_1, \xi_2)$ and $\eta = \begin{pmatrix} {}^t\eta_1 \\ {}^t\eta_2 \end{pmatrix}$ are $N \times 2$ and $2 \times N$ matrices and ξ_i and η_i are fermionic N -vectors. x_0 is a number. The commutation relation is defined by

$$[(X, x_0), (Y, y_0)] = ([X, Y], k \int \frac{dz}{2\pi i} \text{str}(X \partial Y)). \quad (2)$$

The dual space $\hat{\mathbf{g}}^*$ of the superalgebra $\hat{\mathbf{g}}$ consists of an element $(J(z), \alpha_0)$, where the scalar product $\langle \cdot, \cdot \rangle$ take the form

$$\langle (J, \alpha_0), (X, x_0) \rangle = \int \frac{dz}{2\pi i} \text{str}(JX) + \alpha_0 x_0. \quad (3)$$

Let us parametrize $J(z)$ as

$$J(z) = \left(\begin{array}{c|c} A(z) & a(z) \\ \hline b(z) & B(z) \end{array} \right), \quad \text{str} J = \text{tr} A - \text{tr} B = 0. \quad (4)$$

where $A(z) \in gl(N)$, $B(z) \in gl(2)$, $a = (a_1(z), a_2(z))$, $b = \begin{pmatrix} {}^tb_1(z) \\ {}^tb_2(z) \end{pmatrix}$. a_i and b_i ($i = 1, 2$) are N -dimensional vectors whose components are fermionic currents. $\hat{\mathbf{g}}$ acts on $\hat{\mathbf{g}}^*$ as an coadjoint action ad^* (or infinitesimal gauge transformation δ) defined by the formula

$$\langle \text{ad}^*(X, x_0)(J, \alpha_0), (Y, y_0) \rangle = -\langle (J, \alpha_0), [(X, x_0), (Y, y_0)] \rangle, \quad (5)$$

$$i.e. \quad \text{ad}^*(X, x_0)(J, \alpha_0) = ([X, J] + \alpha_0 k \partial X, 0). \quad (6)$$

From (1) and (6), the gauge transformation $\delta J = [X, J] + \alpha_0 k \partial X$ becomes

$$\delta J = \left(\frac{[X_1, A] - a\eta + \xi b + \alpha_0 k \partial X_1}{-B\eta + \eta A - bX_1 + X_2 b + \alpha_0 k \partial \eta} \middle| \frac{-A\xi + \xi B - aX_2 + X_1 a + \alpha_0 k \partial \xi}{[X_2, B] - b\xi + \eta a + \alpha_0 k \partial X_2} \right). \quad (7)$$

Next we impose the constraint on $\hat{\mathbf{g}}^*$. In the present paper, we shall consider the constraint for the $sl(2)$ even subalgebra of $\mathbf{g}$ and keep the other even affine algebra symmetry. We may choose the following constraint set $\mathcal{M}$ (or the set of levels), which consists of the elements of the form $(J(z), 1)$

$$J(z) = \left(\frac{A(z)}{b(z)} \middle| \frac{a(z)}{B(z)} \right), \quad B(z) = \begin{pmatrix} * & 1 \\ * & * \end{pmatrix}. \quad (8)$$

Let $\mathcal{N}$ be the space of gauge transformations which preserve this constraint. The space $\mathcal{N}$ is determined by the condition $\delta B_{12} = 0$ for arbitrary $J(z) \in \mathcal{M}$. From the formula (7), we get $(X_2)_{12} = 0$ and $\xi_2 = 0$ and $\eta_1 = 0$. Namely $\mathcal{N}$ can be expressed as $\mathcal{N} = \mathcal{N}_{u(N)} \mathcal{N}_0$, where $\mathcal{N}_{u(N)}$ is the $u(N)$ gauge transformation group and $\mathcal{N}_0$ consists of the elements e^X with

$$X = \begin{pmatrix} 0 & * & 0 \\ 0 & 0 & 0 \\ * & * & 0 \end{pmatrix}. \quad (9)$$

Let us define the reduced phase space by $\mathcal{F} = \mathcal{M}/\mathcal{N}_0$. We can take the following two typical gauges. One is the ‘‘Drinfeld-Sokolov’’ gauge:

$$J_{DS}(z) = \left(\frac{W}{0} \middle| \frac{G}{0} \frac{0}{1} \right), \quad W = J + \frac{U}{N} 1_N, \quad (10)$$

where J denotes the $sl(N)$ Kac-Moody current at level k and U is the $u(1)$ current, 1_N is the $N \times N$ unit matrix. The currents W generate the $u(N)$ Kac-Moody algebra. The other is the ‘‘diagonal’’ gauge

$$J_{diag}(z) = \left(\frac{A}{\bar{\chi}} \middle| \frac{0}{p_1} \frac{\chi}{1} \right), \quad A = \hat{J} + \frac{p_1 + p_2}{N} 1_N. \quad (11)$$

Here $\hat{J}$ is the $sl(N)$ current at level $\hat{k}$. A Poisson bracket structure on the reduced phase space $\mathcal{F}$ is determined by the coadjoint action preserving the gauge. This method is

known Polyakov's "soldering" procedure [4]. To do this, it is convenient to decompose the gauge parameter by taking $X = \Lambda_1 + \Lambda_2 + \Lambda_3$ where

$$\Lambda_1 = \left(\begin{array}{c|c} \epsilon_1 & 0 \\ \hline 0 & 0 \end{array} \right), \quad \Lambda_2 = \epsilon \left(\begin{array}{c|c} 1_N & 0 \\ \hline 0 & \frac{N}{2} 1_2 \end{array} \right), \quad \Lambda_3 = \left(\begin{array}{c|c} 0 & \xi \\ \hline \eta & \epsilon_2 \end{array} \right), \quad (12)$$

and $\epsilon_1 \in sl(N)$, $\epsilon_2 = \begin{pmatrix} \epsilon_{11} & \epsilon_{12} \\ \epsilon_{21} & -\epsilon_{11} \end{pmatrix} \in sl(2)$. Λ_1 and Λ_2 correspond to the $sl(N)$ Kac-Moody symmetry and the $u(1)$ symmetry, respectively. The parameter Λ_3 belongs to the $sl(2)$ and fermionic part. From the conditions preserving the Drinfeld-Sokolov gauge:

$$\delta a_2 = \delta^t b_1 = 0, \quad \delta B_{11} = \delta B_{12} = 0, \quad (13)$$

we can express the gauge parameters ξ_1 , η_2 , ϵ_{11} and ϵ_{21} in terms of ξ_2 , η_1 , ϵ and ϵ_{12} . Therefore the gauge transformations on the reduced phase space can be expressed by using these parameters:

$$\begin{aligned} \delta U &= Nk\partial\epsilon + {}^t\eta_1 G - {}^t\bar{G}\xi_2, \\ \delta J &= [\epsilon_1, J] + k\partial\epsilon_1 - G^t\eta_1 + \xi_2 {}^t\bar{G} - \frac{1}{N}({}^t\eta_1 G - {}^t\bar{G}\xi_2)1_N \\ \delta G &= -k^2\partial^2\xi_2 + k(2W - U)\partial\xi_2 + \{W(U - W) + k\partial(W - U) + T\}\xi_2 \\ &\quad + \frac{3k}{2}G\partial\epsilon_{12} + \left\{ \left(\frac{1}{2}U - W\right)G + k\partial G \right\} \epsilon_{12} - \left(\frac{N}{2} - 1\right)\epsilon G + \epsilon_1 G, \\ \delta {}^t\bar{G} &= k^2\partial^2 {}^t\eta_1 + k\partial {}^t\eta_1(2W - U) + {}^t\eta_1 \{-T + W(W - U) + k\partial W\} \\ &\quad + \frac{3k}{2} {}^t\bar{G}\partial\epsilon_{12} + \left\{ {}^t\bar{G}(W - \frac{U}{2}) + k\partial {}^t\bar{G} \right\} \epsilon_{12} - {}^t\bar{G}\epsilon_1 + \left(\frac{N}{2} - 1\right) {}^t\bar{G}\epsilon, \\ \delta T &= \frac{N}{2}k^2\partial^2\epsilon - \frac{N}{2}kU\partial\epsilon \\ &\quad - \frac{k^3}{2}\partial^3\epsilon_{12} + \left(2kT + \frac{k}{2}U^2 - k^2\partial U\right)\partial\epsilon_{12} + \left(k\partial T + \frac{k}{2}U\partial U - \frac{k^2}{2}\partial^2 U\right)\epsilon_{12} \\ &\quad + {}^t\eta_1 \{(W - U)G + k\partial G\} + 2k\partial {}^t\eta_1 G + {}^t\bar{G}(U - W)\xi_2 + k {}^t\bar{G}\partial\xi_2. \end{aligned} \quad (14)$$

Note that T itself does not behave as energy-momentum tensor. In order to get the correct form, we must deform T by a differential polynomial of U , so we define $\tilde{T} = T + \frac{1}{4}U^2 - \frac{k}{2}\partial U$.

From (14) we can show that $\tilde{T}$ behaves as

$$\delta \tilde{T} = {}^t\eta_1 \left\{ \left(W - \frac{U}{2}\right)G + \frac{k}{2}\partial G \right\} + \frac{3k}{2}\partial {}^t\eta_1 G + \left\{ {}^t\bar{G}\left(\frac{U}{2} - W\right) + \frac{k}{2}\partial {}^t\bar{G} \right\} \xi_2 + \frac{3k}{2} {}^t\bar{G}\partial\xi_2$$

$$-\frac{k^3}{2}\partial^3\epsilon_{12} + 2k\tilde{T}\partial\epsilon_{12} + k\partial\tilde{T}\epsilon_{12}. \quad (15)$$

By expressing the infinitesimal gauge transformation δ as

$$\delta = \int \frac{dz}{2\pi i} \text{str} \left(\begin{array}{c|cc} W & G & 0 \\ \hline 0 & 0 & 1 \\ \hline {}^t\bar{G} & \tilde{T} & U \end{array} \right) \left(\begin{array}{c|cc} \epsilon_1 + \epsilon_{1N} & 0 & \xi_2 \\ \hline {}^t\eta_1 & \frac{N}{2}\epsilon & \epsilon_{12} \\ \hline 0 & 0 & \frac{N}{2}\epsilon \end{array} \right), \quad (16)$$

we find the operator product expansions for $U, J, G, \bar{G}$ and $\tilde{T}$,

$$\begin{aligned} \tilde{T}(z)\tilde{T}(w) &= \frac{3k^3}{(z-w)^4} + \frac{-2k\tilde{T}}{(z-w)^2} + \frac{-k\partial\tilde{T}}{z-w} + \dots, \\ \tilde{T}(z)G_i(w) &= \frac{\frac{-3k}{2}G_i(w)}{(z-w)^2} + \frac{J_{ij}G_j + (\frac{1}{N} - \frac{1}{2})UG_i - k\partial G_i}{z-w} + \dots, \\ \tilde{T}(z)\bar{G}_i(w) &= \frac{\frac{-3k}{2}\bar{G}_i(w)}{(z-w)^2} + \frac{\bar{G}_jJ_{ji} - (\frac{1}{N} - \frac{1}{2})U\bar{G}_i - k\partial\bar{G}_i}{z-w} + \dots, \\ G_i(z)\bar{G}_j(w) &= \frac{-2k^2\delta_{ij}}{(z-w)^3} + \frac{k(U\delta_{ij} - 2W_{ij})}{(z-w)^2} + \frac{T\delta_{ij} - W_{ik}W_{kj} + W_{ij}U - k\partial W_{ij}}{z-w} + \dots, \\ J_{kl}(z)G_i(w) &= \frac{\delta_{il}G_k - \frac{1}{N}G_i\delta_{kl}}{z-w} + \dots, \quad J_{kl}(z)\bar{G}_i(w) = \frac{-\delta_{ik}\bar{G}_l + \frac{1}{N}\bar{G}_i\delta_{kl}}{z-w} + \dots, \\ J_{ij}(z)J_{kl}(w) &= \frac{k(\delta_{il}\delta_{jk} - \frac{1}{N}\delta_{ij}\delta_{kl})}{(z-w)^2} + \frac{\delta_{jk}J_{il} - \delta_{il}J_{kj}}{z-w} + \dots, \quad U(z)U(w) = \frac{\frac{-2Nk}{N-2}}{(z-w)^2} + \dots, \\ U(z)G_i(w) &= \frac{-G_i(w)}{z-w} + \dots, \quad U(z)\bar{G}_i(w) = \frac{\bar{G}_i(w)}{z-w} + \dots. \end{aligned} \quad (17)$$

The total energy-momentum tensor T_{total} is defined by adding the Sugawara form of energy-momentum tensors of the $sl(N)$ and $u(1)$ Kac-Moody algebras;

$$T_{total}(z) = \frac{1}{-k}\tilde{T}(z) + T_{sl(N)}(z) + T_{u(1)}(z), \quad (18)$$

where $T_{sl(N)} = \frac{1}{2k}\text{tr}J^2$ and $T_{u(1)} = -\frac{N-2}{4Nk}U^2$. The supercurrents $G_i(z)$ and $\bar{G}_i(z)$ have conformal weight $3/2$ with respect to $T_{total}(z)$. These operator product expansions are nothing but those of the classical $u(N)$ extended superconformal algebra. In the case of $N = 2$ the $u(1)$ current decouples from the algebra and we get the $N = 4$ $sl(2)$ ESA.

Next we discuss the Miura transformation by considering the system of linear differential equations [3]:

$$(k\partial - J(z))v(z) = 0, \quad v(z) = \begin{pmatrix} v_0 \\ v_1 \\ v_2 \end{pmatrix}, \quad (19)$$

where $J \in \mathcal{M}$ and v_0 is a N -vector. Under the gauge transformation

$$J(z) \rightarrow n(z)J(z)n(z)^{-1} + k(\partial n(z))n(z)^{-1}, \quad v \rightarrow n(z)v, \quad n(z) \in \mathcal{N}_0, \quad (20)$$

the component v_1 is gauge invariant. Therefore we can write the system of linear differential equations into the gauge-invariant higher (pseudo-)differential operator acting on the component v_1 . In the Drinfeld-Sokolov gauge, the differential equations becomes

$$k\partial v_0 = Wv_0 + Gv_1, \quad k\partial v_1 = v_2, \quad k\partial v_2 = {}^t\bar{G}v_0 + Tv_1 + Uv_2. \quad (21)$$

By solving these equations in terms of v_1 we get

$$\left\{ (k\partial)^2 - Uk\partial - T - {}^t\bar{G}(k\partial - W)^{-1}G \right\} v_1 = 0. \quad (22)$$

Here we define the inverse of $(k\partial - W)$ by

$$(k\partial - W)^{-1} = (k\partial)^{-1} \left\{ 1 + \sum_{i=1}^{\infty} \left(W(k\partial)^{-1} \right)^i \right\}. \quad (23)$$

In the “diagonal” gauge, from the equations

$$k\partial v_0 = Av_0 + \chi v_2, \quad k\partial v_1 = {}^t\bar{\chi}v_0 + p_1v_1 + v_2, \quad k\partial v_2 = p_2v_2, \quad (24)$$

we get

$$\left\{ (k\partial - p_2)(k\partial - p_1) - (k\partial - p_2){}^t\bar{\chi}(k\partial - A + \chi{}^t\bar{\chi})^{-1}\chi(k\partial - p_1) \right\} v_1 = 0. \quad (25)$$

Comparing these two equations (22) and (25), we get the free field representation of the $u(N)$ extended superconformal algebra:

$$\begin{aligned} U &= p_1 + p_2 + {}^t\bar{\chi}\chi, & J &= \hat{J} - \chi{}^t\bar{\chi} - \frac{{}^t\bar{\chi}\chi}{N}1_N, \\ T &= -p_1p_2 + k\partial p_1 + (p_1 + p_2){}^t\bar{\chi}\chi - k(\partial{}^t\bar{\chi})\chi - {}^t\bar{\chi}W\chi, \\ G &= p_1\chi + k\partial\chi - W\chi, & {}^t\bar{G} &= p_2{}^t\bar{\chi} - k\partial{}^t\bar{\chi} - {}^t\bar{\chi}W. \end{aligned} \quad (26)$$

The level $\hat{k}$ of the $sl(N)$ Kac-Moody current $\hat{J}$ is equal to $k - 1$. From eq. (18) the total energy momentum tensor becomes

$$T_{total} = -\frac{1}{2}(\partial\varphi)^2 - \frac{1}{2}(\partial\phi)^2 - \sqrt{\frac{k}{2}}\partial^2\varphi + \frac{1}{2}\left((\partial{}^t\bar{\chi})\chi - {}^t\bar{\chi}\partial\chi\right) + \frac{1}{2k}\text{tr}J^2, \quad (27)$$

where $p_1 - p_2 = \sqrt{2k}\partial\varphi$ and $p_1 + p_2 = \sqrt{\frac{2Nk}{N-2}}\partial\phi$.

In a similar way, we may construct the classical $so(N)$ extended superconformal algebra from the affine Lie superalgebra $osp(N|2)^{(1)}$. An element of $\mathfrak{g} = osp(N|2)$ is given by [12]

$$X = \left(\begin{array}{c|c} A & a \\ \hline b & B \end{array} \right), \quad a = (a_1, a_2), \quad b = \begin{pmatrix} -{}^r a_2 \\ {}^r a_1 \end{pmatrix}, \quad (28)$$

where $A \in so(N)$, $B \in C_1 = A_1$ and a_i ($i = 1, 2$) are N -dimensional fermionic vectors. Here we define a $m \times n$ matrix ${}^r A$ of a $n \times m$ matrix $A = (a_{i,j})$ by ${}^r A = (a_{m+1-j, n+1-i})$. We consider the same type of constraints $\mathcal{M}$ as (8). The gauge group $\mathcal{N}$ which preserve the constraints $\mathcal{M}$ is generated by (9). The soldering procedure can be done by replacing ${}^t \bar{G}$ by ${}^r G$, ${}^t \bar{\chi}$ by $-{}^r \chi$ and p_1 (p_2) by p ($-p$) in the $sl(N|2)$ case, we show only the results

$$\begin{aligned} \delta W &= [\epsilon_1, W] + k\partial\epsilon_1 + G^r \xi_2 + \xi_2 {}^r G \\ \delta G &= -k^2 \partial^2 \xi_2 + kW\partial\xi_2 + \{T - WW + k\partial W\} \xi_2 + \frac{3k}{2} G\partial\epsilon_{12} + (k\partial G - WG) \epsilon_{12} + \epsilon_1 G \\ \delta T &= \frac{-k^3}{2} \partial^3 \epsilon_{12} + 2kT\partial\epsilon_{12} + k\partial T \epsilon_{12} - {}^r \xi_2 (k\partial G + 2WG) - 3k\partial {}^r \xi_2 G. \end{aligned} \quad (29)$$

Here we parametrize the gauge parameters by $\Lambda = \Lambda_1 + \Lambda_3$ in (12) and put η equal to $\begin{pmatrix} -{}^r \xi_2 \\ {}^r \xi_1 \end{pmatrix}$. The Miura transformation ² is given by

$$(k\partial)^2 - T - {}^r G(k\partial - W)^{-1}G = (k\partial + p)(k\partial - p) - (k\partial + p){}^t \bar{\chi}(k\partial - A + \chi {}^r \chi)^{-1}\chi(k\partial - p). \quad (32)$$

Therefore the free field representation of the classical $so(N)$ extended superconformal algebras is expressed as

$$W = \hat{J} - \chi {}^r \chi, \quad G = \sqrt{\frac{k}{2}} \partial\varphi\chi + k\partial\chi - W\chi,$$

²We note that the Miura transformation proposed in refs. [13] ($sl(2|2)$ case) and [11] ($so(N)$ case) is apparently different from the present one. However if we define the path-ordered exponential ‘‘Pexp’’ satisfying

$$\partial \text{Pexp}\left(\int^z W(z')dz'\right) = \text{Pexp}\left(\int^z W(z')dz'\right)W(z), \quad (30)$$

both Miura transformations are shown to be equivalent by virtue of the formula

$$(k\partial - W)^{-1} = \text{Pexp}\left(-\int^z W(z')dz'\right)\partial^{-1}\text{Pexp}\left(\int^z W(z')dz'\right). \quad (31)$$

$$T_{total} = -\frac{1}{2}(\partial\varphi)^2 - \sqrt{\frac{k}{2}}\partial^2\varphi - {}^r\chi\partial\chi + \frac{1}{2k}\text{tr}J^2, \quad (33)$$

where $\hat{J}$ is the $so(N)$ current in the diagonal gauge and the total energy-momentum tensor T_{total} is defined by $-T/k + \text{tr}W^2/(2k)$.

We proceed to the exceptional Lie superalgebra $D(2|1;\alpha)$. The algebra contains even subalgebras $A_1 \oplus A_1 \oplus A_1$ and has rank three. Let α_i ($i = 1, 2, 3$) be simple roots of $D(2|1;\alpha)$ with the inner products $\alpha_1^2 = 0$, $\alpha_2^2 = -2\gamma$, $\alpha_3^2 = -2(1-\gamma)$, $\alpha_1 \cdot \alpha_2 = \gamma$, $\alpha_1 \cdot \alpha_3 = 1-\gamma$ and $\alpha_2 \cdot \alpha_3 = 0$. By constructing the Cartan matrix, we find that the parameter α is given by the relation $\alpha = \gamma/(1-\gamma)$. The positive even roots are α_2 , α_3 and $2\alpha_1 + \alpha_2 + \alpha_3$, which generate A_1 even subalgebras. The odd positive roots are α_1 , $\alpha_1 + \alpha_2$, $\alpha_1 + \alpha_3$ and $\alpha_1 + \alpha_2 + \alpha_3$. For $\alpha = 1$ the hamiltonian reduction can be done by using an explicit 6×6 matrix representation because $D(2|1;\alpha = 1) = osp(4|2)$. But for generic α , there is no such representation. Hence we express the Kac-Moody current as

$$J(z) = \sum_{\alpha \in \Delta_{\bar{0}}} J_{\alpha} e_{\alpha} + \sum_{\alpha \in \Delta_{\bar{1}}} j_{\alpha} e_{\alpha} + \sum_{i=1}^3 H^i h^i, \quad (34)$$

and a gauge parameter $X(z)$ as

$$X(z) = \sum_{\alpha \in \Delta_{\bar{0}}} \epsilon_{\alpha} e_{\alpha} + \sum_{\alpha \in \Delta_{\bar{1}}} \xi_{\alpha} e_{\alpha} + \sum_{i=1}^3 \epsilon^i h^i. \quad (35)$$

Here $\{e_{\alpha}, h^i\}$ are the Cartan-Weyl basis and $\Delta_{\bar{0}}$ ($\Delta_{\bar{1}}$) is the set of even (odd) roots. We denote the currents $J_{\pm\alpha_2}$, $\alpha_1 \cdot H$ as $J_{\pm 2}$, H_1 , etc. for simplicity. Now we consider the constraint on an A_1 subalgebra $\{J_{1123}, J_{-1123}, H_{1123}\}$ by imposing the condition $J_{1123}(z) = 1$. On the reduced phase space we may take the Drinfeld Sokolov gauge

$$\begin{aligned} J_{-1123}(z) &= T(z), \quad H_{1123}(z) = 0, \quad j_{-1}(z) = G_1(z), \\ j_{-12}(z) &= G_{12}(z), \quad j_{-13}(z) = G_{13}(z), \quad j_{-123}(z) = G_{123}(z), \end{aligned} \quad (36)$$

By using the structure constants of the algebra, we can calculate the gauge transformation on the reduced phase space. For Kac-Moody currents the results are

$$\delta H_2 = 2\gamma J_2 \epsilon_{-2} - 2\gamma J_{-2} \epsilon_2 + k\partial(\alpha_2 \cdot \epsilon) - \gamma G_1 \xi_1 + \gamma G_{12} \xi_{12} - \gamma G_{13} \xi_{13} + \gamma G_{123} \xi_{123},$$

$$\begin{aligned}
\delta J_2 &= \gamma G_{13} \xi_{123} - \gamma G_1 \xi_{12} - H_2 \epsilon_2 + k \partial \epsilon_2 + (\alpha_2 \cdot \epsilon) J_2, \\
\delta J_{-2} &= \gamma G_{123} \xi_{13} - \gamma G_{12} \xi_1 + H_2 \epsilon_2 + k \partial \epsilon_{-2} - (\alpha_2 \cdot \epsilon) J_{-2}.
\end{aligned} \tag{37}$$

The transformations of the currents $J_{\pm 3}$ and $\alpha_3 \cdot H$ can be obtained by replacing indices 2 (3) by 3 (2) and γ by $1 - \gamma$. The supercurrents and the energy-momentum tensor transform as

$$\begin{aligned}
\delta G_1 &= J_2 \xi_{-12} + J_3 \xi_{-13} - T \xi_{123} + k \partial \xi_{-1} - \frac{1}{2} (H_2 + H_3) \xi_{-1} \\
&\quad - G_{12} \epsilon_2 - G_{13} \epsilon_3 - (\alpha_1 \cdot \epsilon) G_1, \\
\delta G_{12} &= -J_3 \xi_{-123} + J_{-2} \xi_{-1} - T \xi_{13} + k \partial \xi_{-12} + \frac{1}{2} (H_2 - H_3) \xi_{-12} \\
&\quad + G_{123} \epsilon_3 - G_1 \epsilon_{-2} - (\alpha_{12} \cdot \epsilon) G_{12}, \\
\delta G_{13} &= -J_2 \xi_{-123} + J_{-3} \xi_{-1} - T \xi_{12} + k \partial \xi_{-13} - \frac{1}{2} (H_2 - H_3) \xi_{-13} \\
&\quad + G_{123} \epsilon_2 - G_1 \epsilon_{-3} - (\alpha_{13} \cdot \epsilon) G_{13}, \\
\delta G_{123} &= -J_{-3} \xi_{-12} - J_{-2} \xi_{-13} - T \xi_1 + k \partial \xi_{-123} + \frac{1}{2} (H_2 + H_3) \xi_{-123} \\
&\quad + G_{13} \epsilon_{-2} + G_{12} \epsilon_{-3} - (\alpha_{123} \cdot \epsilon) G_{123}, \\
\delta T &= -(\alpha_{1123} \cdot \epsilon) T + k \partial \epsilon_{-1123} + G_{123} \xi_{-1} + G_{13} \xi_{-12} + G_{12} \xi_{-13} + G_1 \xi_{-123}, \tag{38}
\end{aligned}$$

where $\alpha_{1123} \cdot \epsilon = -k \partial \epsilon_{1123}$ and

$$\begin{aligned}
\xi_{-123} &= -J_{-2} \xi_{12} - J_{-3} \xi_{13} + G_{123} \epsilon_{1123} + k \partial \xi_1 + \frac{1}{2} (H_2 + H_3) \xi_1, \\
\xi_{-13} &= -J_2 \xi_1 + J_{-3} \xi_{123} + G_{13} \epsilon_{1123} + k \partial \xi_{12} - \frac{1}{2} (H_2 - H_3) \xi_{12}, \\
\xi_{-12} &= -J_3 \xi_1 + J_{-2} \xi_{123} + G_{12} \epsilon_{1123} + k \partial \xi_{13} - \frac{1}{2} (-H_2 + H_3) \xi_{13}, \\
\xi_{-1} &= J_2 \xi_{13} + J_3 \xi_{12} + G_1 \epsilon_{1123} + k \partial \xi_{123} - \frac{1}{2} (H_2 + H_3) \xi_{123}, \\
\epsilon_{-1123} &= \epsilon_{1123} T - \frac{1}{2} (G_1 \xi_1 + G_{12} \xi_{12} + G_{13} \xi_{13} + G_{123} \xi_{123}) - \frac{1}{2} k^2 \partial^2 \epsilon_{1123}. \tag{39}
\end{aligned}$$

These relations are those of the non-linear $N = 4$ doubly ESA $\tilde{A}_\gamma$ [14] [15] obtained by factorizing fermions from the linear $N = 4$ ESA A_γ [16]. Generators of $\tilde{A}_\gamma$ are two $sl(2)$ affine Lie algebras $\{J_2, J_{-2}, H_2\}$ at level $k\gamma$, $\{J_3, J_{-3}, H_3\}$ at level $k(1 - \gamma)$, four

supercurrents G_1 , G_{12} , G_{13} and G_{123} and the total energy-momentum tensor $T_{total} = -T/k + T_{Sugawara}$.

One can also calculate the extended superconformal algebras with B_3 and G_2 Kac-Moody symmetries from the hamiltonian reduction of affine Lie superalgebras associated with exceptional Lie superalgebras $F(4)$ and $G(3)$, respectively. In the $F(4)$ case ³, we have eight supercurrents which belong to the $spin(7)$ representation of the B_3 . The G_2 ESA associated with $G(3)$ has seven supercurrents which belong to the 7-dimensional fundamental representation of G_2 . The constructions of these algebras are straightforward by using the matrix (pseudo)representations [18]. From the viewpoint of Lie superalgebra, these complete the classification of extended superconformal algebras. It seems quite interesting to investigate the geometrical meaning of these algebras, because $spin(7)$ and G_2 appear in the context of the classification of restricted holonomy of non-symmetric spaces [19].

It is possible to extend the present analysis to the case of Lie superalgebras $sl(M|N)$ or $osp(M|2N)$, which include an even subalgebra $sl(N)$ or $sp(2N)$. In the former case we get a $Wsl(N)$ minimal model coupled to $gl(M)$ Wess-Zumino-Novikov-Witten (WZNW) model with $2M$ fermionic currents with a conformal weight $(N+1)/2$. In the latter case we get a C_N Toda field theory coupled to the $SO(M)$ WZNW model with N fermionic current with weight $(N+1)/2$.

In a forthcoming paper we shall discuss the quantum hamiltonian reduction and free field representation of extended superconformal algebras. A Feigin-Fuchs type representation enables us to analyse the degenerate representation of the algebra and the computation of correlation functions using screening operators.

We would like to thank Jens Lyng Petersen for useful discussions.

³Recently Fradkin and Linetsky [17] has constructed a quantum version of this algebra by using a non Lie superalgebraic approach.

References

- [1] A.B. Zamolodchikov, Theor. Math. Phys. **63** (1985) 1205.
- [2] V.G. Drinfeld and V.V. Sokolov, J. Sov. Math. **30** (1984) 1975.
- [3] A. A. Belavin, in “Quantum String Theory” Proceedings in Physics vol. **31** (Springer Verlag, Berlin ,1989); M. Bershadsky and H. Ooguri, Commun. Math. Phys.**126** (1989) 429; J. Balog, L. Fehér, P. Forgács, L. O’Raifeartaigh and A. Wipf, Ann. Phys. **203** (1990) 76.
- [4] A.M. Polyakov, Int. J. Mod. Phys. **A5** (1990) 833.
- [5] M.V. Saveliev, Mod. Phys. Lett. **A5** (1990) 2223; M. Bershadsky, Commun. Math. Phys.**139** (1991)71; I. Bakas and D.A. Deprieux, Mod. Phys. Lett. **A6** (1991) 1561; M.F.de Groot, T.J. Hollowood and L.J. Miramontes, preprint IASSNS-HEP-91/19.
- [6] L. Fehér, L. O’Raifeartaigh, P. Ruelle and I. Tsutsui and A. Wipf, Ann. Phys. **213** (1992) 1; preprint DIAS-STP-91-29; P. Bowcock and G.M.T. Watts, preprint EFI 91-63.
- [7] F.A. Bais, T.Tjin and P. van Driel, Nucl. Phys. **B357** (1991) 632; T. Tjin and P.van Driel, preprint ITFA-91-04; J. Fuchs, Phys. Lett. **B262** (1991) 249; L. Romans, Nucl. Phys. **BB357** (1991) 549.
- [8] J. Evans and T. Hollowood, Nucl. Phys. **B352** (1991) 723; S. Komata, K. Mohri and H. Nohara, Nucl. Phys. **B359** (1991) 168; T. Inami and K.-I. Izawa, Phys. Lett. **B255** 521.
- [9] V.G. Knizhnik, Theor. Math. Phys. **66** (1986) 68; M. Bershadsky, Phys. Lett. **B174** (1986) 285.

- [10] B.A. Kupershmidt, Phys. Lett. **A109** (1985) (1985) 417; T.G. Khovanova, Theor. Math. Phys. **72** (1987) 899; M. Bershadsky and H. Ooguri, Phys. Lett. **B229** (1989) 374.
- [11] P. Mathieu, Phys. Lett. **B218** (1989) 185.
- [12] V.G. Kac, Adv. Math. **26** (1977) 8; Lect. Notes in Math. vol. **676**, p.597 (Springer-Verlag, Berlin, 1978).
- [13] M. Chaichian and P.P. Kulish, Phys. Lett. **B183** (1987) 169.
- [14] P. Goddard and A. Schwimmer, Phys. Lett. **B214** (1988) 209.
- [15] J.L. Petersen and A. Taormina, Nucl. Phys. **B331** (1990) 556; **B333** (1990) 833.
- [16] A. Sevrin, W. Troost and A. van Proeyen, Phys. Lett. **B208** (1988) 477.
- [17] E.S. Fradkin and V.Ya. Linetsky, ICTP preprint IC/91/348.
- [18] B.S. DeWitt and P. van Nieuwenhuizen, J. Math. Phys. **23** (1982) 1953.
- [19] M. Berger, Bull. Soc. Math. France, **83** (1955) 279; R.L. Bryant, Lect. Notes in Math., vol. **1111**, p. 269 (Springer-Verlag, Berlin, 1985).

Table 1: classification of extended superconformal algebras

Lie superalgebra	even subalgebra	extended superconformal algebra
$A(n 1) \ (n \geq 2)$	$A_n \oplus A_1 \oplus u(1)$	$N = 2(n+1) \ u(n+1)$ ESA
$A(1 1)$	$A_1 \oplus A_1$	$N = 4 \ sl(2)$ ESA
$A(1 0) = C(2)$	$A_1 \oplus u(1)$	$N = 2$ superconformal algebra
$B(n 1) \ (n \geq 1)$	$B_n \oplus C_1$	$N = 2n+1 \ so(2n+1)$ ESA
$D(n 1) \ (n \geq 2)$	$D_n \oplus C_1$	$N = 2n \ so(2n)$ ESA
$D(2 1; \alpha)$	$A_1 \oplus A_1 \oplus A_1$	$N = 4 \ sl(2) \oplus sl(2)$ ESA
$F(4)$	$B_3 \oplus A_1$	$N = 8 \ spin(7)$ ESA
$G(3)$	$G_2 \oplus A_1$	$N = 7 \ G_2$ ESA